

THE ON-Q* SOLUTION: EFFECTIVE NON-NARCOTIC PAIN RELIEF.¹⁻⁴ ACHIEVE A HIGHER LEVEL OF PATIENT SATISFACTION.^{1,3,5-7}

MANAGING PERIOPERATIVE PAIN IS A CHALLENGE^{9,10}

AN OPTIMAL PLAN CAN IMPROVE PAIN MANAGEMENT

Implement a proven pain management protocol to deliver consistent quality of care^{10,11}

Incorporate continuous nerve blocks as part of a multimodal approach to improve pain control and reduce opioid utilization^{10,11}

Help patients achieve a rapid recovery with reduced length of stay^{1,4,11,12}

Reduce ER visits and readmissions due to pain and opioid-related adverse events^{8,9}

Effectively manage pain beyond discharge improving patient satisfaction^{3,4,7,13,14}

Have expert support for patients and physicians

EFFECTIVE NON-NARCOTIC PAIN RELIEF TO HELP GET YOUR PATIENTS BACK TO NORMAL FASTER

By reducing pain and complications, ON-Q* may help you protect your bundled payment for orthopedic procedures and beyond, while delivering better patient experiences, improving outcomes and reducing opioid utilization.^{1-4,6,7,15,16}

BETTER OUTCOMES^{1,2,4,17,18}

Well-managed pain without narcotics results in fewer complications,^{2-4,6}

Patients will experience fast functional recovery for earlier discharge^{1,2,4,17,18}

SATISFIED PATIENTS^{1,3,5,7}

Improves patient postoperative experiences, which potentially increases HCAHPS scores^{1,3,5-7}

The likelihood of discharging directly home, enhancing patient comfort during recovery^{2,6,7,18}

REDUCES COST OF CARE^{1,2,4,15,17,18}

Potentially reduces length-of-stay and need for skilled nursing facilities^{1,2,4,15,17,18}


Reduces complications that could impact bundled payments, such as ER visits and readmissions^{3,18}

THE ON-Q* SOLUTION

As facilities seek to reduce narcotic use, many have begun using nerve blocks to provide short-term pain relief to patients. One option is a single-shot nerve block, which typically wears off within 24 hours^{19,20}. Alternatively, continuous peripheral nerve blocks (CPNB) deliver significantly better and customizable pain relief beyond 72 hours.^{19,21}

ON-Q* is a disposable pump that automatically and continuously delivers a customizable flow of local anesthetic to a patient's surgical site or in close proximity to nerves.²¹

- Indicated for significantly lower narcotic use and better pain relief than narcotics alone.²¹
- Helps patients feel comfortable while they efficiently move through the recovery process in the hospital and at home.^{1,2,4,6,7,12}
- Offers a seamless transition out of the hospital, as the pump may be sent home with the patient.⁷



ON-Q* IS ON CALL

Avanos delivers best-in-class training and support to help your program succeed.

| | |
|---|---|
| <p>EXPERT IMPLEMENTATION ASSISTANCE</p> | <p>ONGOING TRAINING</p> |
| <p>Over 10 years of experience working with leading institutions to develop successful regional anesthesia programs</p> | <p>Avanos trains over 2000 clinicians annually to stay current with techniques and keep workflow efficient</p> |
| <p>ROUND-THE-CLOCK RESPONSIVENESS</p> | <p>EXTENSIVE LOCAL SUPPORT</p> |
| <p>With nursing-staffed 24/7 hotline for both patients and clinicians—provides reassurance while reducing burden on the care team</p> | <p>Coordinated ON-Q* representatives and clinical experts ensure uninterrupted personalized coverage and continuing success</p> |
| <p>HELP NAVIGATING BUNDLED PAYMENT CHALLENGES</p> | |
| <p>Consult with your team to understand the changing healthcare economic landscape</p> | |

ON-Q* PAIN RELIEF SYSTEM: EFFECTIVE PAIN MANAGEMENT THAT LASTS BEYOND DISCHARGE^{1,4,7}

A total program for keeping pain management from compromising your outcomes and patient satisfaction.^{1,2,5,6,17}

| | | |
|--|----------|--|
| <p>BETTER OUTCOMES TO IMPROVE PATIENT SATISFACTION WITH FEWER COMPLICATIONS AND EARLIER DISCHARGE^{1-7,17,18}</p> | <p>+</p> | <p>REDUCING COST OF CARE MAY POSITIVELY AFFECT YOUR PROGRAM'S REIMBURSEMENT^{1,4,5,9,17,18}</p> |
|--|----------|--|



1. Beaussier M, El'Ayoubi H, Schiffer E, Rollin M, Parc Y, Mazoit J, et al. Continuous preperitoneal infusion of ropivacaine provides effective analgesia and accelerates recovery after colorectal surgery. *Anesthesiology*. 2007;107:461-8.
2. Gómez-Cardero P, Rodríguez-Merchán EC. Postoperative analgesia in TKA: ropivacaine continuous intraarticular infusion. *Clin Orthop Relat Res*. 2010;468(5):1242-1247.
3. Malik T, Mass D, Cohn S. Postoperative analgesia in a prolonged continuous interscalene block versus single-shot block in outpatient arthroscopic rotator cuff repair: a prospective randomized study. *J Arthroscop Rel Surg* 2016: 1-7.
4. Forastiere E, Sofra M, Giannarelli D, Fabrizi L, Simone G. Effectiveness of continuous wound infusion of 0.5% ropivacaine by ON-Q pain relief system for postoperative pain management after open nephrectomy. *Br J Anaesth*. 2008;101(6):841-847.
5. Heller L, Kowalski A, Wei C, Butler C. Prospective, randomized, double-blind trial of local anesthetic infusion and intravenous narcotic patient-controlled anesthesia pump for pain management after free TRAM flap breast reconstruction. *Plast Reconstr Surg*. 2008;122:1010-18.
6. White P, Issioui T, Skrivaneck G, Early J, Wakefield C. The use of continuous popliteal sciatic nerve block after surgery involving the foot and ankle: does it improve the quality of recovery? *Anesth Analg* 2003; 97:1303-9.
7. Visoiu M, Joy LN, Grudziak JS, Chelly JE. The effectiveness of ambulatory continuous peripheral nerve blocks for postoperative pain management in children and adolescents. *Paediatr Anaesth*. 2014;24:1141-1148
8. Oderda GM, Gan TJ, Johnson BH, Robinson SB. Effect of opioid-related adverse events on outcomes in selected surgical patients. *J Pain Palliat Care Pharmacother*. 2013;27:62-70.
9. Kessler ER, Shah M, Gruschkus SK, Raju A. Cost and quality implications of opioid-based postsurgical pain control using administrative claims data from a large health system: opioid-related adverse events and their impact on clinical and economic outcomes. *Pharmacotherapy*. 2013;33(4):383-391.
10. Chou R, Gordon DB, de Leon-Casasola OA, Rosenberg JM, Bickler S, Brennan T, et al. Management of postoperative pain: A clinical practice guideline from the American Pain Society, the American Society of Regional Anesthesia and Pain Medicine, and the American Society of Anesthesiologists' Committee on Regional Anesthesia, Executive Committee, and Administrative Council. *J Pain*. 2016;17(2):131-157.
11. Auyong D, Allen C†, Pahang J, Clabeaux J, MacDonald K, Hanson N. Reduced length of hospitalization in primary total knee arthroplasty patients using an updated enhanced recovery after orthopedic surgery (ERAS) pathway. *J Arthroplasty*. 2015;30:1705-9.
12. Mudumbai SC, Kim TE, Howard SK, Workman JJ, Giori N, Woolson S, et al. Continuous adductor canal blocks are superior to continuous femoral nerve blocks in promoting early ambulation after TKA. *Clin Orthop Relat Res*. 2014;472(5):1377-1383.
13. Liu S, Richman J, Thirlby R, Wu C. Efficacy of continuous wound catheters delivering local anesthetic for postoperative analgesia: a quantitative and qualitative systematic review of randomized controlled trials. *J Am Coll Surg*. 2006;203:914-32.
14. Hachem F, Canar J, Fullam F, Gallan A, Hohmann S, Johnson C. The relationships between HCAHPS communication and discharge satisfaction items and hospital readmissions. *Patient Experience Journal*. 2014;1:71-7.
15. Dowling R, Thielmeier K, Ghaly A, Barber D, Boice T, Dine A. Improved pain control after cardiac surgery: results of a randomized, double-blind, clinical trial. *J Thorac Cardiovasc Surg*. 2003;126(5):1271-1278. †
16. Goyal N, McKenzie J, Sharkey PF, Parvizi J, Hozack WJ, Austin MS. The 2012 Chitranjan Ranawat award: intraarticular analgesia after TKA reduces pain: a randomized, double-blinded, placebo-controlled, prospective study. *Clin Orthop Relat Res*. 2013;471(1):64-75.
17. White PF, Rawal S, Latham P, Markowitz S, Issioui T, Chi L, Dellaria S, et al. Use of a continuous local anesthetic infusion for pain management after median sternotomy. *Anesthesiology*. 2003;99(4):918-923.
18. Dulaney-Cripe E, Hadaway S, Bauman R, Trame C, Sillaman B, Laughlin R. A continuous infusion fascia iliac compartment block in hip fracture patients: a pilot study. *J Clin Med Res* 2012; 45-48.
19. Bingham A, Fu R, Horn J, Abrahams M. Continuous peripheral nerve block compared with single-injection peripheral nerve block: a systematic review and meta-analysis of randomized controlled trials. *Reg Anesth Pain Med*. 2012;37:583-94.
20. Abdallah FW, Halpern SH, Aoyama K, Brull R. Will the real benefits of single-shot interscalene block please stand up? A systematic review and meta-analysis. *Anesth Analg*. 2015;120(5):1114-1129 21. The ON-Q* Pain Relief System. Instructions for use. Avanos Medical, Inc.
21. The ON-Q* Pain Relief System. Instructions for use. Avanos Medical, Inc. † Dr. Auyong has a consulting/speaking/financial relationship with Avanos Medical, Inc.

There are inherent risks in all medical devices. Please refer to the product labeling for **Indications, Cautions, Warnings and Contraindications**. Failure to follow the product labeling could directly impact patient safety. Physician is responsible for prescribing and administering medications per instructions provided by the drug manufacturer. Refer to www.avanospainmanagement.com for additional product safety Technical Bulletins.